


LIST OF CLAIMS

1. (Currently Amended) Aminooxy-cyclodextrin derivatives compounds of the formula 1:



wherein

CD is a mono- or polydeoxy α -, β - or γ -cyclodextrin, carrying in its 6-, 3- and/or 2-position the aminooxy function containing group (X-Y-ONH₂), and optionally carrying further substituents different from (X-Y-ONH₂) in their 6-, 3- and/or 2-positions, and wherein Y is a linker group between the aminooxy group and the mono- or polydeoxy-CD-group,

X is a functional group or an atom necessary to connect the linker Y and the deoxy CD group, or Y is a direct bond when X is a direct bond, and

n is greater than or equal to 1, but less than or equal to 18, 21 or 24 for α -, β - or γ -cyclodextrin, respectively, as well as the aminooxy protected derivatives thereof.

2. (Currently Amended) The derivative compound according to claim 1, wherein Y and X are both direct bonds.

3. (Currently Amended) The derivative compound according to claim 1 or 2, wherein one or more of the primary hydroxyl groups at

a 6-position of α -, β - or γ -CD are substituted with a X-Y-ONH₂ fragment, wherein X and Y have the meaning of claim 1.

4. (Currently Amended) The derivative compound according to claim 1, wherein Y is a linear or branched alkylene, alkenylene with one or more double ^{bonds} bounds which may be either isolated or conjugated, alkynylene with one or more triple bonds which may be either isolated or conjugated, or arylene or arylalkylene fragments where aryl may be substituted or not substituted, whereby the alkylene, alkenylene and alkynylene fragments may be linear or branched, and one or more of the chain members (methylene groups) may be replaced by -NH-, -O-, -S-, -S-S-, -C(O)NH-, -C(O)O-, -OP(O)(OH)O-, -S(O)-, SO₂-, or -CHR-, where R is alkyl, aryl, -OR', -NH₂, -NHR', -NR'₂, -OH, -COOH, or -ONH₂ groups and where R' is alkyl, aryl, or acyl.

5. (Currently Amended) The derivative compound according to claims 1 or 4, wherein X is selected from the group consisting of -O-, -S-, -NH-, -NR"-, -OCO-, -NH-O-, =NO-, -NHC(O)-, -OP(O)(OH), and -R"C=NO-, where R" is linear or branched lower alkyl.


6. (Currently Amended) The derivative compound according to claim 4, wherein Y is alkylene containing 2-12 C-atoms, wherein one

or more of the chain members may be replaced by -NH-, -O-, -S-, -C(O)NH-, -C(O)O-, or CHR₁ wherein R₁ is methyl, ethyl or propyl and X is -O-, -S-, -NH-, -OC(O)-, or -NH-C(O)-.

7. (Currently Amended) The derivative compound according to claim 1, wherein one or more of the hydroxyl groups at 6-, 3-, and/or 2-position(s) are substituted with a group selected from the group consisting of H₂N-, HS-, -COOH, alkoxy-, aryloxy-, and acyloxy, and wherein said alkoxy-, aryloxy-, and acyloxy- can contain H₂N-, HS-, or -COOH in their structure, side chain or aromatic ring.


8. (Currently Amended/Withdrawn from Consideration) A method for preparing the derivative compound of claim 1 of formula 1, wherein X is an oxygen atom, comprising the steps of:

a) alkylating a cyclodextrin of formula (3) at one or more of the positions 6, 3, and/or 2 containing a hydroxyl group,




wherein R_2 , R_3 , and R_4 are hydrogen or substituents selected from the group consisting of alkyl, aryl and acyl, and wherein said substituents' functional groups, if they exist, are protected whenever necessary,

with a compound according to formula (3'):




wherein W is $-OC_2H_5$ or $-CH_3$, m and Y are as defined in claim 1, and Z is a reactive group, and optionally protecting group(s) is/are removed, or

b) alkylating a cyclodextrin derivative compound of formula (4)




wherein R'_2 , R'_3 , R'_4 are hydroxy or activated groups selected from the group consisting of tosyl, mesyl, halogen, ester and epoxy/~~or~~ said substituent being in a protected form if necessary, whereby the cyclodextrin derivative compound contains at least one of said activated groups

with the compound of formula (4')



wherein X and Y are as in claims 1, 4 or 5, and W is $-\text{OC}_2\text{H}_5$ or $-\text{CH}_3$, and protecting group(s) is/are removed if necessary, or

(c) reacting a cyclodextrin ~~derivative~~ compound of formula (5)



wherein at least one of the groups R''_2 , R''_3 , and R''_4 are thiol-, amino-, carboxy-, or alkoxy-, aryloxy- or acyloxy groups which contain at least one thiol-, amino-, carboxy- group, or their derivative, and the remaining functional groups are hydroxyl groups or they have the meaning described in claim 7 for the substituents, and exist, if necessary, in a protected form, with an appropriate aminooxy protected substituted hydroxylamine according to formula (3'), after which the protecting group(s) are removed, or

(d) reacting a cyclodextrine ~~derivative~~ compound of formula (5), which contains one or more of keto or aldehyde groups, with bisaminooxy alkanes of formula (5')



wherein t is 2-12, and wherein one of the methylene groups can be substituted with oxygen or sulfur atom, or wherein -NH- or -S-S- groups, and a protecting group is removed if necessary.

9. (Canceled)

Delete

10. (Currently Amended/Withdrawn from Consideration) An oxime created from any one of the aminooxy-CDs of claim 1 by reacting said aminooxy-CDs with synthetic or natural aldehydes or ketones.

Delete

11. (Currently Amended/Withdrawn from Consideration) ~~Derivatives~~ Compounds of nucleotide or nucleoside pyrimidines or purines with aminooxy-CDs, wherein said aminooxy group is linked to the heterocyclic ring of said pyrimidines or purines.

12. (Currently Amended) The aminooxy-cyclodextrin ~~derivatives~~ compounds of claim 1, wherein the aminooxy protected ~~derivative~~ compound is ethoxy-ethylidene or acetone oxime derivatives thereof.


13. (Currently Amended) The ~~derivatives~~ compounds according to claim 4, where the alkylene, alkenylene, and alkynylene fragments contain 2 to 12 c-atoms in the chain.

14. (Currently Amended) The ~~derivatives~~ compounds of claim 7, where the alkoxy is a C₁-C₆ alkoxy, the aryloxy is phenyloxy, benzyloxy or tolyloxy, and the acyloxy originates from C₁-C₆ carboxyl or benzoic acids.

~~Delete~~ 15. (Currently Amended/Withdrawn from Consideration) The ~~derivatives~~ compounds of claim 11, wherein the aminoxy group is linked to the heterocyclic ring through pyrimidine C-4 and purine C-6, and wherein pyrimidine and purine are cytosine or adenine, or their derivatives.


16. (Currently Amended/Withdrawn from Consideration) A method for preparing the ~~derivative~~ compound of formula 1 of claim 3, wherein X is an oxygen atom, comprising the steps of: a) alkylating a cyclodextrin of formula (3), at one or more of the positions 6, 3, and/or 2 containing a hydroxyl group, wherein R₂, R₃, and R₄ are hydrogen or substituents selected from the group consisting of alkyl, aryl, and acyl, and wherein said substituents' functional groups, if they exist, are protected whenever necessary,

with a compound according to formula (3'):




wherein W is $-\text{OC}_2\text{H}_5$ or $-\text{CH}_3$, m and Y are as defined in claim 3, and Z is a reactive group, and optionally protecting group(s) is/are removed, or

b) alkylating a cyclodextrin derivative compound of formula (4)




wherein R'_2 , R'_3 , R'_4 are hydroxy or activated groups selected from the group consisting of tosyl, mesyl, halogen, ester and epoxy or said substituent being in a protected form if necessary, whereby the cyclodextrin derivative compound contains at least one of said activated groups

with the compound of formula (4')



wherein X and Y are as in claims 1, 4 or 5 and W is $-\text{OC}_2\text{H}_5$ or $-\text{CH}_3$, and protecting group(s) is/are removed if necessary, or

(c) reacting a cyclodextrin derivative compound of formula (5)



wherein at least one of the groups R''_2 , R''_3 , and R''_4 are thiol-, amino-, carboxy-, or alkoxy-, aryloxy-, or acyloxy groups which contain at least one thiol-, amino-, carboxy-group, or their derivative, and the remaining functional groups are hydroxyl groups or they have the meaning described in claim 7 for the substituents, and exist, if necessary, in a protected form, modified with an appropriate aminooxy protected substituted hydroxylamine according to formula (3'), after which the protecting group(s) are removed, or

(d) reacting a cyclodextrin derivative compound of formula (5), which contains one or more of keto or aldehyde groups, with bisaminooxy alkanes of formula (5')



wherein t is 2-12, and wherein one of the methylene groups can be substituted with oxygen or sulfur atom, or wherein -NH- or -S-S- groups, and a protecting group is removed if necessary.

17. (New) The aminoxy-cyclodextrin compounds of the formula 1 as recited in claim 1, wherein Y is a linker group represented by a linear or branched alkylene, alkenylene with one or more double bonds which may be either isolated or conjugated, alkynylene with one or more triple bonds which may be either isolated or conjugated, or arylene or arylalkylene fragments where aryl may be substituted or not substituted, whereby the alkylene, alkenylene and alkynylene fragments may be linear or branched, and one or more of the chain members (methylene groups) may be replaced by -NH-, -O-, -S-, -S-S-, -C(O)NH-, -C(O)O-, -OP(O)(OH)O-, -S(O)-, SO₂-, or -CHR-, where R is alkyl, aryl, -OR', -NH₂, -NHR', -NR'₂, -OH, -COOH, or -ONH₂ groups and where R' is alkyl, aryl, or acyl.

18. (New) The aminoxy-cyclodextrin compounds of the formula 1 as recited in claim 17, wherein the amino protective group is ethoxy-ethylidene or acetone oxime derivatives thereof.

19. (New) The aminoxy-cyclodextrin compounds of the formula 1 as recited in claim 18, wherein X is selected from the group consisting of -O-, -S-, -NH-, -NR"-, -OCO-, -NH-O-, =NO-, -NHC(O)-, -OP(O)(OH), and -R"C=NO-, where R" is linear or branched lower alkyl.